Methodist Health System

How to Provide a Clean Catch Mid-Void Urine Sample – Female

Important: It is important to follow the instructions below exactly, to properly clean the bacteria from the skin and urinary opening. A dirty specimen may result in the need for a second test, or worse yet, may be the cause of a wrong diagnosis, unnecessary medicine, or the wrong medicine being given to you.

Instructions for Females:

- 1. Wash your hands with soap and water, then rinse and dry.
- 2. Open all the towelettes and place them on a clean surface (i.e, a cloth or paper towel.)
- 3. Place towelettes and specimen container within easy reach.
- 4. Remove all undergarments.
- 5. Sit comfortably on the toilet seat and position one leg to the side as far as possible.
- 6. With one hand, spread the labia folds and keep them spread during the cleansing and collection of the urine specimen.
- 7. **WASH** the area from which the urine is passed. Use the three soap towelettes. Using one towelette at a time, begin to wash, gently wiping only from front to back between the folds of skin. Discard that towelette. Repeat the front and back wash process, using one towelette at a time.
- 8. **RINSE**. Rinse with one sterile saline towelette. Use the front to back motion.
- 9. **VOID** a small amount of urine into the toilet; then collect a portion of the urine in the specimen container. The cup should be held to avoid contact with skin and clothing. Keep the fingers away from the rim and inner surface of the container.

Questions About Your Test:

If you have any questions about the test, call the doctor that ordered the test or the Pathology Center, 402.354.4541 or 888.432.8980.

Reviewed 11/2022 Nebraska Methodist Health System. All rights reserved. All content provided herein is for educational, informational and guidance purposes only and is not intended to serve as a substitute for individualized professional medical advice, diagnosis, or treatment. Nothing contained herein establishes or shall be used to establish the legal definition of the "standard of care." The content should be used as guidance and, since it may not be universally applied to all patients in all situations, healthcare professionals should use the content along with independent judgment and on a case by case basis.