









<p align="center">Midstream Collection Kit Lawson Ordering - 200149</p>	<p align="center">Individual Cup Lawson Ordering - 194111</p>	<p align="center">Adapter for Foley Catheter Urine Transfer Lawson Ordering - 47513</p>
 <ul style="list-style-type: none"> Instruct patient not to remove yellow seal. Remove seal and insert tubes into specially designed lid for withdrawal. Discard blue lid in sharps. Label/send tubes to laboratory. <p>Use for routine midstream collections.</p>	 <ul style="list-style-type: none"> Can be ordered individually if specific unit does not perform Urinalysis and Culture together. See below for individual vacutainer tube ordering. 	 <ul style="list-style-type: none"> Attach device to foley catheter. Select/insert appropriate tubes for withdrawal. Discard device in sharps. Label/send tubes to laboratory.

<p align="center">Urinalysis Preservative Tube Lawson Ordering – 79335</p>	<p align="center">Urine Culture Preservative Tube Lawson Ordering - 79332</p>	<p align="center">Urine Tube Non – Additive Lawson Ordering - 58469</p>
 <p>Volume: 7 mL (min) – 8 mL Used for:</p> <ul style="list-style-type: none"> Urinalysis 	 <p>Volume: 3 mL (min) – 4 mL Used for:</p> <ul style="list-style-type: none"> Urine Culture 	 <p>Volume: 6 mL (preferred full) Used for:</p> <ul style="list-style-type: none"> Urine Chemistry Urine Pregnancy Urine Toxicology Miscellaneous Sendout
<p align="center">Routinely collected <u>together</u> for Urinalysis with the possibility of added Urine Culture</p>		<p align="center">*See list below for ≥6 mL volume urine samples.</p>

<p align="center">Urine Cytology Collections</p>	<p align="center"><i>Affix patient label containing a minimum of 2 forms of identification to each specimen.</i></p>	
 <p>Due to larger volume requirement for Urine Cytology (50 mL), current collection will continue.</p> <ul style="list-style-type: none"> Collect in specimen cup Hand-Deliver to Laboratory 	<ul style="list-style-type: none"> Urine vacutainer tubes are designed to enhance specimen integrity <i>and</i> minimize spill risk when transported through the pneumatic tube system. If “Add On” testing is a possibility, it may be best to send all three tubes. <p align="center">Questions? Contact the Clinical Laboratory at (720) 848-4401</p>	<p align="center">* Urine Testing, Volumes Greater than 6 mL</p> <p align="center">15 ml Minimum (3 tubes)</p> <ul style="list-style-type: none"> Protein Electrophoresis ED Abuse Drug Screen Urine Drugs of Abuse Drug of Abuse with Quantitation (15 mL)
		<p align="center">10 ml Minimum (2 tubes)</p> <ul style="list-style-type: none"> Iodine Random Urine (10 mL) Mucopolysaccharides (10 mL)