

Longs Peak Hospital - Clinical Laboratory

Stool Collection Guide

GI PCR Panel Plus Stool Culture	Ova and Parasite Exam (O&P)	C. difficile toxin	Fecal Occult Blood
Para-Pak C85 The second secon	EXINC-PVA FOR PORMACING and The Proposed and Thomas and	SALUE AT YCLICK* ST-7000 Full Nome: DO B:	Lisean Prompt* reason to the second of the s
Collection Container – Stool culture kit:	Collection Container – O&P kit	Collection Container - Sterile cup	Collection Container - Sterile cup
 1 vial with Cary Blair (orange or green lid) 1 empty vial (clear lid) 1 set of collection instructions 1 ziplock bag Do not use if pink indicator fluid color fades or turns yellow. 	 1 Zinc-PVA Fixative vial (gray lid) And 1 Formalin vial (pink lid) Must send both vials 	5 – 10 mL fresh stool Specimens in media (such as Cary Blair) are not accepted Formed stools are not accepted	Fresh stool applied to occult blood card
Fill to line. Do not overfill! Use a tongue depressor or collection spoon attached to lid Vigorously shake fluid vials.	Fill to line. Do not overfill! Use a tongue depressor or collection spoon attached to lid Vigorously shake fluid vials.		Scrape surface of fecal sample with the sampling probe, completely covering the grooved portion. Apply to occult blood card
Stability:	Stability:	Stability:	Stability:
Preserved stool 72 hours ambient Fresh stool, 30 minutes ambient	Indefinite / ambient	Deliver to lab immediately Refrigerate if offsite ≤ 24 hours	10 days ambient