

190.25 - Alpha-fetoprotein

Other Names/Abbreviations

AFP

Description

Alpha-fetoprotein (AFP) is a polysaccharide found in some carcinomas. It is effective as a biochemical marker for monitoring the response of certain malignancies to therapy.

HCPCS Codes (Alphanumeric, CPT[©] AMA)

Code	Description
82105	Alpha-fetoprotein; serum
	· · · · · · · · · · · · · · · · · · ·

ICD-10-CM Codes Covered by Medicare Program

The ICD-10-CM codes in the table below can be viewed on CMS' website as part of Downloads: Lab Code List, at http://www.cms.gov/Medicare/Coverage/CoverageGenInfo/LabNCDsICD10.html

Code	Description
A52.74	Syphilis of liver and other viscera
B18.0	Chronic viral hepatitis B with delta-agent
B18.1	Chronic viral hepatitis B without delta-agent
B18.2	Chronic viral hepatitis C
B66.1	Clonorchiasis
B66.3	Fascioliasis
C22.0	Liver cell carcinoma
C22.1	Intrahepatic bile duct carcinoma
C22.2	Hepatoblastoma
C22.3	Angiosarcoma of liver
C22.4	Other sarcomas of liver
C22.7	Other specified carcinomas of liver
C22.8	Malignant neoplasm of liver, primary, unspecified as to type
C22.9	Malignant neoplasm of liver, not specified as primary or secondary
C38.1	Malignant neoplasm of anterior mediastinum
C38.2	Malignant neoplasm of posterior mediastinum
C38.3	Malignant neoplasm of mediastinum, part unspecified
C38.8	Malignant neoplasm of overlapping sites of heart, mediastinum and pleura

NCD 190.25



Code	Description
C44.1321	Sebaceous cell carcinoma of skin of right upper eyelid, including canthus
C44.1322	Sebaceous cell carcinoma of skin of right lower eyelid, including canthus
C44.1391	Sebaceous cell carcinoma of skin of left upper eyelid, including canthus
C44.1392	Sebaceous cell carcinoma of skin of left lower eyelid, including canthus
C56.1	Malignant neoplasm of right ovary
C56.2	Malignant neoplasm of left ovary
C56.3	Malignant neoplasm of bilateral ovaries
C56.9	Malignant neoplasm of unspecified ovary
C62.00	Malignant neoplasm of unspecified undescended testis
C62.01	Malignant neoplasm of undescended right testis
C62.02	Malignant neoplasm of undescended left testis
C62.10	Malignant neoplasm of unspecified descended testis
C62.11	Malignant neoplasm of descended right testis
C62.12	Malignant neoplasm of descended left testis
C62.90	Malignant neoplasm of unspecified testis, unspecified whether descended or undescended
C62.91	Malignant neoplasm of right testis, unspecified whether descended or undescended
C62.92	Malignant neoplasm of left testis, unspecified whether descended or undescended
C78.1	Secondary malignant neoplasm of mediastinum
C78.7	Secondary malignant neoplasm of liver and intrahepatic bile duct
C79.60	Secondary malignant neoplasm of unspecified ovary
C79.61	Secondary malignant neoplasm of right ovary
C79.62	Secondary malignant neoplasm of left ovary
C79.63	Secondary malignant neoplasm of bilateral ovaries
C79.82	Secondary malignant neoplasm of genital organs
C7A.00	Malignant carcinoid tumor of unspecified site
C7A.090	Malignant carcinoid tumor of the bronchus and lung
C7A.091	Malignant carcinoid tumor of the thymus
C7A.092	Malignant carcinoid tumor of the stomach
C7A.093	Malignant carcinoid tumor of the kidney
C7A.094	Malignant carcinoid tumor of the foregut, unspecified
C7A.095	Malignant carcinoid tumor of the midgut, unspecified
C7A.096	Malignant carcinoid tumor of the hindgut, unspecified
C7A.098	Malignant carcinoid tumors of other sites
C7B.00	Secondary carcinoid tumors, unspecified site

NCD 190.25



Code	Description
C7B.01	Secondary carcinoid tumors of distant lymph nodes
C7B.02	Secondary carcinoid tumors of liver
C7B.03	Secondary carcinoid tumors of bone
C7B.04	Secondary carcinoid tumors of peritoneum
C7B.09	Secondary carcinoid tumors of other sites
C7B.1	Secondary Merkel cell carcinoma
C7B.8	Other secondary neuroendocrine tumors
D13.4	Benign neoplasm of liver
D13.5	Benign neoplasm of extrahepatic bile ducts
D37.6	Neoplasm of uncertain behavior of liver, gallbladder and bile ducts
D64.0	Hereditary sideroblastic anemia
D64.1	Secondary sideroblastic anemia due to disease
D64.2	Secondary sideroblastic anemia due to drugs and toxins
D64.3	Other sideroblastic anemias
D81.810	Biotinidase deficiency
D81.82	Activated Phosphoinositide 3-kinase Delta Syndrome [APDS]
D84.1	Defects in the complement system
E78.2	Mixed hyperlipidemia
E83.00	Disorder of copper metabolism, unspecified
E83.01	Wilson's disease
E83.09	Other disorders of copper metabolism
E83.10	Disorder of iron metabolism, unspecified
E83.110	Hereditary hemochromatosis
E83.111	Hemochromatosis due to repeated red blood cell transfusions
E83.118	Other hemochromatosis
E83.119	Hemochromatosis, unspecified
E83.19	Other disorders of iron metabolism
E84.19	Cystic fibrosis with other intestinal manifestations
E84.9	Cystic fibrosis, unspecified
E88.01	Alpha-1-antitrypsin deficiency
E88.02	Plasminogen deficiency
G89.3	Neoplasm related pain (acute) (chronic)
125.84	Coronary atherosclerosis due to calcified coronary lesion
174.01	Saddle embolus of abdominal aorta

NCD 190.25



Code	Description
174.09	Other arterial embolism and thrombosis of abdominal aorta
174.10	Embolism and thrombosis of unspecified parts of aorta
174.19	Embolism and thrombosis of other parts of aorta
J98.59	Other diseases of mediastinum, not elsewhere classified
K70.2	Alcoholic fibrosis and sclerosis of liver
K70.30	Alcoholic cirrhosis of liver without ascites
K70.31	Alcoholic cirrhosis of liver with ascites
K73.0	Chronic persistent hepatitis, not elsewhere classified
K73.1	Chronic lobular hepatitis, not elsewhere classified
K73.2	Chronic active hepatitis, not elsewhere classified
K73.8	Other chronic hepatitis, not elsewhere classified
K73.9	Chronic hepatitis, unspecified
K74.00	Hepatic fibrosis, unspecified
K74.01	Hepatic fibrosis, early fibrosis
K74.02	Hepatic fibrosis, advanced fibrosis
K74.60	Unspecified cirrhosis of liver
K74.69	Other cirrhosis of liver
K75.4	Autoimmune hepatitis
K76.81	Hepatopulmonary syndrome
K76.82	Hepatic encephalopathy
N44.1	Cyst of tunica albuginea testis
N44.2	Benign cyst of testis
N44.8	Other noninflammatory disorders of the testis
N50.3	Cyst of epididymis
N50.811	Right testicular pain
N50.812	Left testicular pain
N50.819	Testicular pain, unspecified
N50.82	Scrotal pain
N50.89	Other specified disorders of the male genital organs
N53.12	Painful ejaculation
N53.8	Other male sexual dysfunction
N53.9	Unspecified male sexual dysfunction
Q53.111	Unilateral intraabdominal testis
Q53.112	Unilateral inguinal testis

NCD 190.25



Code	Description
Q53.13	Unilateral high scrotal testis
Q53.211	Bilateral intraabdominal testes
Q53.212	Bilateral inguinal testes
Q53.23	Bilateral high scrotal testes
R39.83	Unilateral non-palpable testicle
R39.84	Bilateral non-palpable testicles
R91.1	Solitary pulmonary nodule
R91.8	Other nonspecific abnormal finding of lung field
R93.1	Abnormal findings on diagnostic imaging of heart and coronary circulation
R93.2	Abnormal findings on diagnostic imaging of liver and biliary tract
R93.5	Abnormal findings on diagnostic imaging of other abdominal regions, including retroperitoneum
R93.811	Abnormal radiologic findings on diagnostic imaging of right testicle
R93.812	Abnormal radiologic findings on diagnostic imaging of left testicle
R93.813	Abnormal radiologic findings on diagnostic imaging of testicles, bilateral
R93.819	Abnormal radiologic findings on diagnostic imaging of unspecified testicle
R93.89	Abnormal findings on diagnostic imaging of other specified body structures
R97.8	Other abnormal tumor markers
Z17.0	Estrogen receptor positive status [ER+]
Z17.1	Estrogen receptor negative status [ER-]
*Z17.21	*Progesterone receptor positive status
*Z17.22	*Progesterone receptor negative status
*Z17.31	*Human epidermal growth factor receptor 2 positive status
*Z17.32	*Human epidermal growth factor receptor 2 negative status
*Z17.410	*Hormone receptor positive with human epidermal growth factor receptor 2 positive status
*Z17.411	*Hormone receptor positive with human epidermal growth factor receptor 2 negative status
*Z17.420	*Hormone receptor negative with human epidermal growth factor receptor 2 positive status
*Z17.421	*Hormone receptor negative with human epidermal growth factor receptor 2 negative status
Z85.05	Personal history of malignant neoplasm of liver
Z85.43	Personal history of malignant neoplasm of ovary
Z85.47	Personal history of malignant neoplasm of testis
Z86.002	Personal history of in-situ neoplasm of other and unspecified genital organs

NCD 190.25



Code	Description
Z86.003	Personal history of in-situ neoplasm of oral cavity, esophagus and stomach
Z86.004	Personal history of in-situ neoplasm of other and unspecified digestive organs

Indications

AFP is useful for the diagnosis of hepatocellular carcinoma in high-risk patients (such as alcoholic cirrhosis, cirrhosis of viral etiology, hemochromatosis, and alpha 1-antitrypsin deficiency) and in separating patients with benign hepatocellular neoplasms or metastases from those with hepatocellular carcinoma and, as a non-specific tumor associated antigen, serves in marking germ cell neoplasms of the testis, ovary, retro peritoneum, and mediastinum.

ICD-10-CM Codes That Do Not Support Medical Necessity

Any ICD-10-CM code not listed in either of the ICD-10-CM covered or non-covered sections.

Sources of Information

Tatsuta M. Yamamura H. Iishi H. Kasugai H. Okuda S.Value of serum alpha-fetoprotein and ferritin in the diagnosis of hepatocellular carcinoma. Oncology. 43(5):306-10, 1986.



190.26 - Carcinoembryonic Antigen

Other Names/Abbreviations

CEA

Description

Carcinoembryonic antigen (CEA) is a protein polysaccharide found in some carcinomas. It is effective as a biochemical marker for monitoring the response of certain malignancies to therapy.

HCPCS Codes (Alphanumeric, CPT[©] AMA)

Code	Description
82378	Carcinoembryonic antigen (CEA)

ICD-10-CM Codes Covered by Medicare Program

The ICD-10-CM codes in the table below can be viewed on CMS' website as part of Downloads: Lab Code List, at http://www.cms.gov/Medicare/Coverage/CoverageGenInfo/LabNCDsICD10.html

Code	Description
C15.3	Malignant neoplasm of upper third of esophagus
C15.4	Malignant neoplasm of middle third of esophagus
C15.5	Malignant neoplasm of lower third of esophagus
C15.8	Malignant neoplasm of overlapping sites of esophagus
C15.9	Malignant neoplasm of esophagus, unspecified
C16.0	Malignant neoplasm of cardia
C16.1	Malignant neoplasm of fundus of stomach
C16.2	Malignant neoplasm of body of stomach
C16.3	Malignant neoplasm of pyloric antrum
C16.4	Malignant neoplasm of pylorus
C16.5	Malignant neoplasm of lesser curvature of stomach, unspecified
C16.6	Malignant neoplasm of greater curvature of stomach, unspecified
C16.8	Malignant neoplasm of overlapping sites of stomach
C16.9	Malignant neoplasm of stomach, unspecified
C17.0	Malignant neoplasm of duodenum
C17.1	Malignant neoplasm of jejunum
C17.2	Malignant neoplasm of ileum
C17.3	Meckel's diverticulum, malignant

NCD 190.26



Code	Description
C17.8	Malignant neoplasm of overlapping sites of small intestine
C17.9	Malignant neoplasm of small intestine, unspecified
C18.0	Malignant neoplasm of cecum
C18.1	Malignant neoplasm of appendix
C18.2	Malignant neoplasm of ascending colon
C18.3	Malignant neoplasm of hepatic flexure
C18.4	Malignant neoplasm of transverse colon
C18.5	Malignant neoplasm of splenic flexure
C18.6	Malignant neoplasm of descending colon
C18.7	Malignant neoplasm of sigmoid colon
C18.8	Malignant neoplasm of overlapping sites of colon
C18.9	Malignant neoplasm of colon, unspecified
C19	Malignant neoplasm of rectosigmoid junction
C20	Malignant neoplasm of rectum
C21.0	Malignant neoplasm of anus, unspecified
C21.1	Malignant neoplasm of anal canal
C21.2	Malignant neoplasm of cloacogenic zone
C21.8	Malignant neoplasm of overlapping sites of rectum, anus and anal canal
C25.0	Malignant neoplasm of head of pancreas
C25.1	Malignant neoplasm of body of pancreas
C25.2	Malignant neoplasm of tail of pancreas
C25.3	Malignant neoplasm of pancreatic duct
C25.4	Malignant neoplasm of endocrine pancreas
C25.7	Malignant neoplasm of other parts of pancreas
C25.8	Malignant neoplasm of overlapping sites of pancreas
C25.9	Malignant neoplasm of pancreas, unspecified
C26.0	Malignant neoplasm of intestinal tract, part unspecified
C33	Malignant neoplasm of trachea
C34.00	Malignant neoplasm of unspecified main bronchus
C34.01	Malignant neoplasm of right main bronchus
C34.02	Malignant neoplasm of left main bronchus
C34.10	Malignant neoplasm of upper lobe, unspecified bronchus or lung
C34.11	Malignant neoplasm of upper lobe, right bronchus or lung
C34.12	Malignant neoplasm of upper lobe, left bronchus or lung

NCD 190.26



Code	Description
C34.2	Malignant neoplasm of middle lobe, bronchus or lung
C34.30	Malignant neoplasm of lower lobe, unspecified bronchus or lung
C34.31	Malignant neoplasm of lower lobe, right bronchus or lung
C34.32	Malignant neoplasm of lower lobe, left bronchus or lung
C34.80	Malignant neoplasm of overlapping sites of unspecified bronchus and lung
C34.81	Malignant neoplasm of overlapping sites of right bronchus and lung
C34.82	Malignant neoplasm of overlapping sites of left bronchus and lung
C34.90	Malignant neoplasm of unspecified part of unspecified bronchus or lung
C34.91	Malignant neoplasm of unspecified part of right bronchus or lung
C34.92	Malignant neoplasm of unspecified part of left bronchus or lung
C44.1321	Sebaceous cell carcinoma of skin of right upper eyelid, including canthus
C44.1322	Sebaceous cell carcinoma of skin of right lower eyelid, including canthus
C44.1391	Sebaceous cell carcinoma of skin of left upper eyelid, including canthus
C44.1392	Sebaceous cell carcinoma of skin of left lower eyelid, including canthus
C50.011	Malignant neoplasm of nipple and areola, right female breast
C50.012	Malignant neoplasm of nipple and areola, left female breast
C50.019	Malignant neoplasm of nipple and areola, unspecified female breast
C50.021	Malignant neoplasm of nipple and areola, right male breast
C50.022	Malignant neoplasm of nipple and areola, left male breast
C50.029	Malignant neoplasm of nipple and areola, unspecified male breast
C50.111	Malignant neoplasm of central portion of right female breast
C50.112	Malignant neoplasm of central portion of left female breast
C50.119	Malignant neoplasm of central portion of unspecified female breast
C50.121	Malignant neoplasm of central portion of right male breast
C50.122	Malignant neoplasm of central portion of left male breast
C50.129	Malignant neoplasm of central portion of unspecified male breast
C50.211	Malignant neoplasm of upper-inner quadrant of right female breast
C50.212	Malignant neoplasm of upper-inner quadrant of left female breast
C50.219	Malignant neoplasm of upper-inner quadrant of unspecified female breast
C50.221	Malignant neoplasm of upper-inner quadrant of right male breast
C50.222	Malignant neoplasm of upper-inner quadrant of left male breast
C50.229	Malignant neoplasm of upper-inner quadrant of unspecified male breast
C50.311	Malignant neoplasm of lower-inner quadrant of right female breast
C50.312	Malignant neoplasm of lower-inner quadrant of left female breast

NCD 190.26



Code	Description
C50.319	Malignant neoplasm of lower-inner quadrant of unspecified female breast
C50.321	Malignant neoplasm of lower-inner quadrant of right male breast
C50.322	Malignant neoplasm of lower-inner quadrant of left male breast
C50.329	Malignant neoplasm of lower-inner quadrant of unspecified male breast
C50.411	Malignant neoplasm of upper-outer quadrant of right female breast
C50.412	Malignant neoplasm of upper-outer quadrant of left female breast
C50.419	Malignant neoplasm of upper-outer quadrant of unspecified female breast
C50.421	Malignant neoplasm of upper-outer quadrant of right male breast
C50.422	Malignant neoplasm of upper-outer quadrant of left male breast
C50.429	Malignant neoplasm of upper-outer quadrant of unspecified male breast
C50.511	Malignant neoplasm of lower-outer quadrant of right female breast
C50.512	Malignant neoplasm of lower-outer quadrant of left female breast
C50.519	Malignant neoplasm of lower-outer quadrant of unspecified female breast
C50.521	Malignant neoplasm of lower-outer quadrant of right male breast
C50.522	Malignant neoplasm of lower-outer quadrant of left male breast
C50.529	Malignant neoplasm of lower-outer quadrant of unspecified male breast
C50.611	Malignant neoplasm of axillary tail of right female breast
C50.612	Malignant neoplasm of axillary tail of left female breast
C50.619	Malignant neoplasm of axillary tail of unspecified female breast
C50.621	Malignant neoplasm of axillary tail of right male breast
C50.622	Malignant neoplasm of axillary tail of left male breast
C50.629	Malignant neoplasm of axillary tail of unspecified male breast
C50.811	Malignant neoplasm of overlapping sites of right female breast
C50.812	Malignant neoplasm of overlapping sites of left female breast
C50.819	Malignant neoplasm of overlapping sites of unspecified female breast
C50.821	Malignant neoplasm of overlapping sites of right male breast
C50.822	Malignant neoplasm of overlapping sites of left male breast
C50.829	Malignant neoplasm of overlapping sites of unspecified male breast
C50.911	Malignant neoplasm of unspecified site of right female breast
C50.912	Malignant neoplasm of unspecified site of left female breast
C50.919	Malignant neoplasm of unspecified site of unspecified female breast
C50.921	Malignant neoplasm of unspecified site of right male breast
C50.922	Malignant neoplasm of unspecified site of left male breast
C50.929	Malignant neoplasm of unspecified site of unspecified male breast

NCD 190.26



Code	Description
C56.1	Malignant neoplasm of right ovary
C56.2	Malignant neoplasm of left ovary
C56.3	Malignant neoplasm of bilateral ovaries
C56.9	Malignant neoplasm of unspecified ovary
C78.00	Secondary malignant neoplasm of unspecified lung
C78.01	Secondary malignant neoplasm of right lung
C78.02	Secondary malignant neoplasm of left lung
C78.4	Secondary malignant neoplasm of small intestine
C78.5	Secondary malignant neoplasm of large intestine and rectum
C7A.00	Malignant carcinoid tumor of unspecified site
C7A.010	Malignant carcinoid tumor of the duodenum
C7A.011	Malignant carcinoid tumor of the jejunum
C7A.012	Malignant carcinoid tumor of the ileum
C7A.019	Malignant carcinoid tumor of the small intestine, unspecified portion
C7A.020	Malignant carcinoid tumor of the appendix
C7A.021	Malignant carcinoid tumor of the cecum
C7A.022	Malignant carcinoid tumor of the ascending colon
C7A.023	Malignant carcinoid tumor of the transverse colon
C7A.024	Malignant carcinoid tumor of the descending colon
C7A.025	Malignant carcinoid tumor of the sigmoid colon
C7A.026	Malignant carcinoid tumor of the rectum
C7A.029	Malignant carcinoid tumor of the large intestine, unspecified portion
C7A.090	Malignant carcinoid tumor of the bronchus and lung
C7A.091	Malignant carcinoid tumor of the thymus
C7A.092	Malignant carcinoid tumor of the stomach
C7A.093	Malignant carcinoid tumor of the kidney
C7A.094	Malignant carcinoid tumor of the foregut, unspecified
C7A.095	Malignant carcinoid tumor of the midgut, unspecified
C7A.096	Malignant carcinoid tumor of the hindgut, unspecified
C7A.098	Malignant carcinoid tumors of other sites
C7B.00	Secondary carcinoid tumors, unspecified site
C7B.01	Secondary carcinoid tumors of distant lymph nodes
C7B.02	Secondary carcinoid tumors of liver
C7B.03	Secondary carcinoid tumors of bone

NCD 190.26



Code	Description
C7B.04	Secondary carcinoid tumors of peritoneum
C7B.09	Secondary carcinoid tumors of other sites
C7B.1	Secondary Merkel cell carcinoma
C7B.8	Other secondary neuroendocrine tumors
D01.0	Carcinoma in situ of colon
D01.1	Carcinoma in situ of rectosigmoid junction
D01.2	Carcinoma in situ of rectum
D01.40	Carcinoma in situ of unspecified part of intestine
D01.49	Carcinoma in situ of other parts of intestine
D01.7	Carcinoma in situ of other specified digestive organs
D01.9	Carcinoma in situ of digestive organ, unspecified
D37.1	Neoplasm of uncertain behavior of stomach
D37.2	Neoplasm of uncertain behavior of small intestine
D37.3	Neoplasm of uncertain behavior of appendix
D37.4	Neoplasm of uncertain behavior of colon
D37.5	Neoplasm of uncertain behavior of rectum
G89.3	Neoplasm related pain (acute) (chronic)
R70.1	Abnormal plasma viscosity
R77.0	Abnormality of albumin
R77.1	Abnormality of globulin
R77.2	Abnormality of alphafetoprotein
R77.8	Other specified abnormalities of plasma proteins
R77.9	Abnormality of plasma protein, unspecified
R78.89	Finding of other specified substances, not normally found in blood
R78.9	Finding of unspecified substance, not normally found in blood
R79.89	Other specified abnormal findings of blood chemistry
R97.0	Elevated carcinoembryonic antigen [CEA]
R97.8	Other abnormal tumor markers
Z08	Encounter for follow-up examination after completed treatment for malignant neoplasm
Z09	Encounter for follow-up examination after completed treatment for conditions other than malignant neoplasm
Z85.00	Personal history of malignant neoplasm of unspecified digestive organ
Z85.038	Personal history of other malignant neoplasm of large intestine
Z85.048	Personal history of other malignant neoplasm of rectum, rectosigmoid junction, and anus

NCD 190.26



Code	Description
Z85.118	Personal history of other malignant neoplasm of bronchus and lung
Z85.3	Personal history of malignant neoplasm of breast
Z85.43	Personal history of malignant neoplasm of ovary
Z86.002	Personal history of in-situ neoplasm of other and unspecified genital organs
Z86.003	Personal history of in-situ neoplasm of oral cavity, esophagus and stomach
Z86.004	Personal history of in-situ neoplasm of other and unspecified digestive organs

Indications

CEA may be medically necessary for follow-up of patients with colorectal carcinoma. It would however only be medically necessary at treatment decision-making points. In some clinical situations (e.g. adenocarcinoma of the lung, small cell carcinoma of the lung, and some gastrointestinal carcinomas) when a more specific marker is not expressed by the tumor, CEA may be a medically necessary alternative marker for monitoring. Preoperative CEA may also be helpful in determining the post-operative adequacy of surgical resection and subsequent medical management. In general, a single tumor marker will suffice in following patients with colorectal carcinoma or other malignancies that express such tumor markers.

In following patients who have had treatment for colorectal carcinoma, ASCO guideline suggests that if resection of liver metastasis would be indicated, it is recommended that post-operative CEA testing be performed every two to three months in patients with initial stage II or stage III disease for at least two years after diagnosis.

For patients with metastatic solid tumors which express CEA, CEA may be measured at the start of the treatment and with subsequent treatment cycles to assess the tumor's response to therapy.

Limitations

Serum CEA determinations are generally not indicated more frequently than once per chemotherapy treatment cycle for patients with metastatic solid tumors which express CEA or every two months post-surgical treatment for patients who have had colorectal carcinoma. However, it may be proper to order the test more frequently in certain situations, for example, when there has been a significant change from prior CEA level or a significant change in patient status which could reflect disease progression or recurrence.

Testing with a diagnosis of an in situ carcinoma is not reasonably done more frequently than once, unless the result is abnormal, in which case the test may be repeated once.

ICD-10-CM Codes That Do Not Support Medical Necessity

Any ICD-10-CM code not listed in either of the ICD-10-CM covered or non-covered sections.

Sources of Information

Journal Clinical Oncol: 14(10:2843-2877), 1996

NCD 190.26

*January 2025 Changes ICD-10-CM Version – Red



Vauthey JN. Dudrick PS. Lind DS. Copeland EM 3rd. Management of recurrent colorectal cancer: another look at carcinoembryonic antigen detected recurrence [see comments]. [Review] Digestive Diseases. 14(1):5©13, 1996 Jan-Feb.

Germ J. The prognostic importance of tumor markers in adenocarcinoma of the gastrointestinal tract. [Review] [38 refs] Current Opinion in Oncology. 9(4):380-7, 1997 Jul.

Bergama chi R. Arnaud JP. Routine compared with nonscheduled follow-up of patients with "curative" surgery for colorectal cancer. Annals of Surgical Oncology. 3(5):464-9, 1996 Sep.

Kim YH. Ajani JA. Ota DM. Lynch P. Roth JA. Value of serial carcinoembryonic antigen levels in patients with respectable adenocarcinoma of the esophagus and stomach Cancer. 75(2):451©6, 1995 Jan 15.



190.28 - Tumor Antigen by Immunoassay CA 125

Description

Immunoassay determinations of the serum levels of certain proteins or carbohydrates serve as tumor markers. When elevated, serum concentration of these markers may reflect tumor size and grade. This policy specifically addresses tumor antigen CA 125.

HCPCS Codes (Alphanumeric, CPT[®] AMA)

Code	Description
86304	Immunoassay for tumor antigen, quantitative, CA 125

ICD-10-CM Codes Covered by Medicare Program

The ICD-10-CM codes in the table below can be viewed on CMS' website as part of Downloads: Lab Code List, at http://www.cms.gov/Medicare/Coverage/CoverageGenInfo/LabNCDsICD10.html

Code	Description
C45.1	Mesothelioma of peritoneum
C48.1	Malignant neoplasm of specified parts of peritoneum
C48.2	Malignant neoplasm of peritoneum, unspecified
C48.8	Malignant neoplasm of overlapping sites of retroperitoneum and peritoneum
C51.8	Malignant neoplasm of overlapping sites of vulva
C53.0	Malignant neoplasm of endocervix
C54.1	Malignant neoplasm of endometrium
C54.2	Malignant neoplasm of myometrium
C54.3	Malignant neoplasm of fundus uteri
C54.9	Malignant neoplasm of corpus uteri, unspecified
C56.1	Malignant neoplasm of right ovary
C56.2	Malignant neoplasm of left ovary
C56.3	Malignant neoplasm of bilateral ovaries
C56.9	Malignant neoplasm of unspecified ovary
C57.00	Malignant neoplasm of unspecified fallopian tube
C57.01	Malignant neoplasm of right fallopian tube
C57.02	Malignant neoplasm of left fallopian tube
C57.4	Malignant neoplasm of uterine adnexa, unspecified
C57.7	Malignant neoplasm of other specified female genital organs
C57.8	Malignant neoplasm of overlapping sites of female genital organs

NCD 190.28



Code	Description
C79.60	Secondary malignant neoplasm of unspecified ovary
C79.61	Secondary malignant neoplasm of right ovary
C79.62	Secondary malignant neoplasm of left ovary
C79.63	Secondary malignant neoplasm of bilateral ovaries
C79.82	Secondary malignant neoplasm of genital organs
D39.0	Neoplasm of uncertain behavior of uterus
D39.10	Neoplasm of uncertain behavior of unspecified ovary
D39.11	Neoplasm of uncertain behavior of right ovary
D39.12	Neoplasm of uncertain behavior of left ovary
D39.2	Neoplasm of uncertain behavior of placenta
D39.8	Neoplasm of uncertain behavior of other specified female genital organs
D39.9	Neoplasm of uncertain behavior of female genital organ, unspecified
G89.3	Neoplasm related pain (acute) (chronic)
M33.03	Juvenile dermatomyositis without myopathy
M33.13	Other dermatomyositis without myopathy
M33.93	Dermatopolymyositis, unspecified without myopathy
R19.01	Right upper quadrant abdominal swelling, mass and lump
R19.02	Left upper quadrant abdominal swelling, mass and lump
R19.03	Right lower quadrant abdominal swelling, mass and lump
R19.04	Left lower quadrant abdominal swelling, mass and lump
R19.09	Other intra-abdominal and pelvic swelling, mass and lump
R97.1	Elevated cancer antigen 125 [CA 125]
R97.8	Other abnormal tumor markers
Z85.41	Personal history of malignant neoplasm of cervix uteri
Z85.42	Personal history of malignant neoplasm of other parts of uterus
Z85.43	Personal history of malignant neoplasm of ovary
Z85.44	Personal history of malignant neoplasm of other female genital organs

Indications

CA 125 is a high molecular weight serum tumor marker elevated in 80% of patients who present with epithelial ovarian carcinoma. It is also elevated in carcinomas of the fallopian tube, endometrium, and endocervix. An elevated level may also be associated with the presence of a malignant mesothelioma or primary peritoneal carcinoma.

NCD 190.28



A CA 125 level may be obtained as part of the initial pre-operative work-up for women presenting with a suspicious pelvic mass to be used as a baseline for purposes of post-operative monitoring. Initial declines in CA 125 after initial surgery and/or chemotherapy for ovarian carcinoma are also measured by obtaining three serum levels during the first month post treatment to determine the patient's CA 125 half-life, which has significant prognostic implications.

The CA 125 levels are again obtained at the completion of chemotherapy as an index of residual disease. Surveillance CA 125 measurements are generally obtained every 3 months for 2 years, every 6 months for the next 3 years, and yearly thereafter. CA 125 levels are also an important indicator of a patient's response to therapy in the presence of advanced or recurrent disease. In this setting, CA 125 levels may be obtained prior to each treatment cycle.

Limitations

These services are not covered for the evaluation of patients with signs or symptoms suggestive of malignancy. The service may be ordered at times necessary to assess either the presence of recurrent disease or the patient's response to treatment with subsequent treatment cycles.

The CA 125 is specifically not covered for aiding in the differential diagnosis of patients with a pelvic mass as the sensitivity and specificity of the test is not sufficient. In general, a single "tumor marker" will suffice in following a patient with one of these malignancies.

ICD-10-CM Codes That Do Not Support Medical Necessity

Any ICD-10-CM code not listed in either of the ICD-10-CM covered or non-covered sections.

Documentation Requirements

Indicated if service request for CA125 is requested more frequently than stipulated.

Sources of Information

Clinical Pancreatic Guideline for the Use of Tumor Markers in Breast and Colorectal Cancer, American Society of Clinical Oncology. J Clin Oncol 14:2843-2877, 1996.

Chan DW, Beveridge RA, Muss H, et al. Use of Triquant BR Radioimmunoassay for Early Detection of Breast Cancer Recurrence in Patients with Stage II and Stage III Disease. J Clin Oncol 1977, 15(6):2322-2328.

NCD 190.28



190.29 - Tumor Antigen by Immunoassay CA 15-3/CA 27.29

Description

Immunoassay determinations of the serum levels of certain proteins or carbohydrates serve as tumor markers. When elevated, serum concentration of markers may reflect tumor size & grade. This policy specifically addresses the following tumor antigens: CA 15-3 and CA 27.29

HCPCS Codes (Alphanumeric, CPT[©] AMA)

Code	Description
86300	Immunoassay for tumor antigen, quantitative; CA 15-3 (27.29)

ICD-10-CM Codes Covered by Medicare Program

The ICD-10-CM codes in the table below can be viewed on CMS' website as part of Downloads: Lab Code List, at

http://www.cms.gov/Medicare/Coverage/CoverageGenInfo/LabNCDsICD10.html

Code	Description
C44.1321	Sebaceous cell carcinoma of skin of right upper eyelid, including canthus
C44.1322	Sebaceous cell carcinoma of skin of right lower eyelid, including canthus
C44.1391	Sebaceous cell carcinoma of skin of left upper eyelid, including canthus
C44.1392	Sebaceous cell carcinoma of skin of left lower eyelid, including canthus
C50.011	Malignant neoplasm of nipple and areola, right female breast
C50.012	Malignant neoplasm of nipple and areola, left female breast
C50.019	Malignant neoplasm of nipple and areola, unspecified female breast
C50.021	Malignant neoplasm of nipple and areola, right male breast
C50.022	Malignant neoplasm of nipple and areola, left male breast
C50.029	Malignant neoplasm of nipple and areola, unspecified male breast
C50.111	Malignant neoplasm of central portion of right female breast
C50.112	Malignant neoplasm of central portion of left female breast
C50.119	Malignant neoplasm of central portion of unspecified female breast
C50.121	Malignant neoplasm of central portion of right male breast
C50.122	Malignant neoplasm of central portion of left male breast
C50.129	Malignant neoplasm of central portion of unspecified male breast
C50.211	Malignant neoplasm of upper-inner quadrant of right female breast
C50.212	Malignant neoplasm of upper-inner quadrant of left female breast
C50.219	Malignant neoplasm of upper-inner quadrant of unspecified female breast
C50.221	Malignant neoplasm of upper-inner quadrant of right male breast

NCD 190.29

*January 2025 Changes ICD-10-CM Version – Red



Code	Description
C50.222	Malignant neoplasm of upper-inner quadrant of left male breast
C50.229	Malignant neoplasm of upper-inner quadrant of unspecified male breast
C50.311	Malignant neoplasm of lower-inner quadrant of right female breast
C50.312	Malignant neoplasm of lower-inner quadrant of left female breast
C50.319	Malignant neoplasm of lower-inner quadrant of unspecified female breast
C50.321	Malignant neoplasm of lower-inner quadrant of right male breast
C50.322	Malignant neoplasm of lower-inner quadrant of left male breast
C50.329	Malignant neoplasm of lower-inner quadrant of unspecified male breast
C50.411	Malignant neoplasm of upper-outer quadrant of right female breast
C50.412	Malignant neoplasm of upper-outer quadrant of left female breast
C50.419	Malignant neoplasm of upper-outer quadrant of unspecified female breast
C50.421	Malignant neoplasm of upper-outer quadrant of right male breast
C50.422	Malignant neoplasm of upper-outer quadrant of left male breast
C50.429	Malignant neoplasm of upper-outer quadrant of unspecified male breast
C50.511	Malignant neoplasm of lower-outer quadrant of right female breast
C50.512	Malignant neoplasm of lower-outer quadrant of left female breast
C50.519	Malignant neoplasm of lower-outer quadrant of unspecified female breast
C50.521	Malignant neoplasm of lower-outer quadrant of right male breast
C50.522	Malignant neoplasm of lower-outer quadrant of left male breast
C50.529	Malignant neoplasm of lower-outer quadrant of unspecified male breast
C50.611	Malignant neoplasm of axillary tail of right female breast
C50.612	Malignant neoplasm of axillary tail of left female breast
C50.619	Malignant neoplasm of axillary tail of unspecified female breast
C50.621	Malignant neoplasm of axillary tail of right male breast
C50.622	Malignant neoplasm of axillary tail of left male breast
C50.629	Malignant neoplasm of axillary tail of unspecified male breast
C50.811	Malignant neoplasm of overlapping sites of right female breast
C50.812	Malignant neoplasm of overlapping sites of left female breast
C50.819	Malignant neoplasm of overlapping sites of unspecified female breast
C50.821	Malignant neoplasm of overlapping sites of right male breast
C50.822	Malignant neoplasm of overlapping sites of left male breast
C50.829	Malignant neoplasm of overlapping sites of unspecified male breast
C50.911	Malignant neoplasm of unspecified site of right female breast
C50.912	Malignant neoplasm of unspecified site of left female breast

NCD 190.29



Code	Description
C50.919	Malignant neoplasm of unspecified site of unspecified female breast
C50.921	Malignant neoplasm of unspecified site of right male breast
C50.922	Malignant neoplasm of unspecified site of left male breast
C50.929	Malignant neoplasm of unspecified site of unspecified male breast
C79.2	Secondary malignant neoplasm of skin
C79.81	Secondary malignant neoplasm of breast
C84.7A	Anaplastic large cell lymphoma, ALK-negative, breast
*C84.7B	*Anaplastic large cell lymphoma, ALK-negative, in remission
G89.3	Neoplasm related pain (acute) (chronic)
R97.8	Other abnormal tumor markers
Z85.3	Personal history of malignant neoplasm of breast
Z86.002	Personal history of in-situ neoplasm of other and unspecified genital organs
Z86.003	Personal history of in-situ neoplasm of oral cavity, esophagus and stomach
Z86.004	Personal history of in-situ neoplasm of other and unspecified digestive organs
Z86.005	Personal history of in-situ neoplasm of middle ear and respiratory system
Z86.006	Personal history of melanoma in-situ
Z86.007	Personal history of in-situ neoplasm of skin

Indications

Multiple tumor markers are available for monitoring the response of certain malignancies to therapy and assessing whether a residual tumor exists post-surgical therapy. CA 15-3 is often medically necessary to aid in the management of patients with breast cancer. Serial testing must be used in conjunction with other clinical methods for monitoring breast cancer. For monitoring, if medically necessary, use consistently either CA 15-3 or CA 27.29, not both. CA 27.29 is equivalent to CA 15-3 in its usage in management of patients with breast cancer.

Limitations

These services are not covered for the evaluation of patients with signs or symptoms suggestive of malignancy. The service may be ordered at times necessary to assess either the presence of recurrent disease or the patient's response to treatment with subsequent treatment cycles.

ICD-10-CM Codes That Do Not Support Medical Necessity

Any ICD-10-CM code not listed in either of the ICD-10-CM covered or non-covered sections.

Sources of Information

Clinical Pancreatic Guideline for the Use of Tumor Markers in Breast and Colorectal Cancer, American Society of Clinical Oncology. J Clin Oncol 14:2843-2877, 1996.

NCD 190.29

*January 2025 Changes ICD-10-CM Version – Red

Fu Associates, Ltd.

January 2025



Chan DW, Beveridge RA, Muss H, et al. Use of Triquant BR Radioimmunoassay for Early Detection of Breast Cancer Recurrence in Patients with Stage II & Stage III Disease. J Clin Oncol 1977, 15(6):2322-2328.

Bone GG, von Mensdorff-Pouilly S, Kenemans P, van Kamp GJ, et al. Clinical and Technical Evaluation of ACS BR Serum Assay of MUC-1 Gene Derived Glycoprotein in Breast Cancer, and Compared with CA15-3 Assays. Clin Chem 1997, 43(4):585-593.

NCD 190.29

Fu Associates, Ltd.



190.30 - Tumor Antigen by Immunoassay CA 19-9

Description

Immunoassay determinations of the serum levels of certain proteins or carbohydrates serve as tumor markers. When elevated, serum concentration of these markers may reflect tumor size and grade. This policy specifically addresses the following tumor antigen: CA19-9.

HCPCS Codes (Alphanumeric, CPT[©] AMA)

Code	Description
86301	Immunoassay for tumor antigen, quantitative; CA 19-9

ICD-10-CM Codes Covered by Medicare Program

The ICD-10-CM codes in the table below can be viewed on CMS' website as part of Downloads: Lab Code List, at http://www.cms.gov/Medicare/Coverage/CoverageGenInfo/LabNCDsICD10.html

Code	Description
C22.1	Intrahepatic bile duct carcinoma
C23	Malignant neoplasm of gallbladder
C24.0	Malignant neoplasm of extrahepatic bile duct
C24.1	Malignant neoplasm of ampulla of Vater
C24.8	Malignant neoplasm of overlapping sites of biliary tract
C24.9	Malignant neoplasm of biliary tract, unspecified
C25.0	Malignant neoplasm of head of pancreas
C25.1	Malignant neoplasm of body of pancreas
C25.2	Malignant neoplasm of tail of pancreas
C25.3	Malignant neoplasm of pancreatic duct
C25.4	Malignant neoplasm of endocrine pancreas
C25.7	Malignant neoplasm of other parts of pancreas
C25.8	Malignant neoplasm of overlapping sites of pancreas
C25.9	Malignant neoplasm of pancreas, unspecified
C78.7	Secondary malignant neoplasm of liver and intrahepatic bile duct
C78.80	Secondary malignant neoplasm of unspecified digestive organ
C78.89	Secondary malignant neoplasm of other digestive organs
D37.6	Neoplasm of uncertain behavior of liver, gallbladder and bile ducts
D37.8	Neoplasm of uncertain behavior of other specified digestive organs
D37.9	Neoplasm of uncertain behavior of digestive organ, unspecified

NCD 190.30



Code	Description
G89.3	Neoplasm related pain (acute) (chronic)
M33.03	Juvenile dermatomyositis without myopathy
M33.13	Other dermatomyositis without myopathy
M33.93	Dermatopolymyositis, unspecified without myopathy
R97.8	Other abnormal tumor markers
Z85.068	Personal history of other malignant neoplasm of small intestine
Z85.07	Personal history of malignant neoplasm of pancreas
Z85.09	Personal history of malignant neoplasm of other digestive organs

Indications

Multiple tumor markers are available for monitoring the response of certain malignancies to therapy and assessing whether residual tumor exists post-surgical therapy.

Levels are useful in following the course of patients with established diagnosis of pancreatic and biliary ductal carcinoma. The test is not indicated for diagnosing these two diseases.

Limitations

These services are not covered for the evaluation of patients with signs or symptoms suggestive of malignancy. The service may be ordered at times necessary to assess either the presence of recurrent disease or the patient's response to treatment with subsequent treatment cycles.

ICD-10-CM Codes That Do Not Support Medical Necessity

Any ICD-10-CM code not listed in either of the ICD-10-CM covered or non-covered sections.

Sources of Information

Clinical Pancreatic Guideline for the Use of Tumor Markers in Breast and Colorectal Cancer, American Society of Clinical Oncology. J Clin Oncol 14:2843-2877, 1996.

Richter JM, Christensen MR, Rustgi AK, and Silverstein MD. The Clinical Utility of the CA19-9 Radioimmunoassay for the Diagnosis of Pancreatic Cancer Presenting as Pain or Weight Loss: A Cost Effective Analysis. Arch Intern Med 1989, 149:2292-2297.

Safi F, SchlosseW, Falkenreck S, et. al. Prognostic Value of CA 19-9 Serum Course in Pancreatic Cancer. Hepaetogastroenterology 1998 Jan-Feb; 45(19):253-9.